C62_Proceeding_ICTAP_A_Conn ection_for_A_Partially.pdf

by Wipsar Sunu Brams Dwandaru

Submission date: 12-Dec-2021 08:20AM (UTC+0700)

Submission ID: 1727830994

File name: C62_Proceeding_ICTAP_A_Connection_for_A_Partially.pdf (741.93K)

Word count: 4315

Character count: 22879



A Connection for a Partially Asymmetric Driven Lattice Gas of a Mean-Field Type to an Exact Lattice Density Functional Via a Diagrammatic Method



TC 1-21

A Connection for a Partially Asymmetric Driven Lattice Gas of a Mean-Field Type to an Exact Lattice Density Functional Via a Diagrammatic Method

Wipsar Sunu Brams Dwandaru^{1,*} and Matthias Schmidt²

¹Physics Education Department, Faculty of Mathematics and Natural Sciences, Universitas Negeri Yogyakarta, Karangmalang Complex, 55281, Yogyakarta, Indonesia

²Theoretische Physik II, Physikalisches Institut Universität Bayreuth, D-95440, Germany

*Email: wipsarian@uny.ac.id

Abstract. A relationship of the totally asymmetric simple exclusion process (TASEP) of particles on a one-dimensional lattice system that interact with site exclusion to an exact density functional has been established in [1]. Here, the connection is extended for a partially asymmetric exclusion process (PASEP) where particles on a one-dimensional lattice system may be driven to the left or to the right with certain (left and right) hopping rates. The mean-field equation for the one-particle density of the PASEP is equivalent to the exact equilibrium density of a lattice fluids mixture with three species. The connection is put forward by associating the particles in the lattice fluids mixture to the particle and the particle movements in the PASEP. The particle densities of the lattice fluids mixture is obtained based on the lattice fundamental measure theory via a diagrammatic method. Omitting one of the particles in the mixture yields back the relationship given in [1].

Keywords: PASEP, hopping rates, lattice fluids mixture. **PACS:** 05

INTRODUCTION

Exclusion processes are a class of driven diffusive systems which are widely used to model non-equilibrium phenomena such as transport processes in biological systems [2,3], biopolymerization [4,5], vehicular traffics [6-11], and performance of wireless network [12]. This model consists of hard-core particles moving (jumping) through a one-dimensional lattice sites with certain dynamical rule and boundary conditions. In this paper in particular, the model being studied is the partially asymmetric simple exclusion process (PASEP). Here, particles may move to the right (left) nearest neighbor site with hopping rate $k_+(k_-)$. If one of the hopping rate is set to zero, e.g. $k_{-} = 0$, then the well known totally asymmetric simple exclusion process (TASEP) [13, 14] is obtained.

A relationship based on a mean-field theory between the TASEP and the equilibrium lattice gas through classical density functional theory is given in [1]. This relationship is obtained by a correspondence between the appropriate particles of the binary mixture to the resting and moving particles of the TASEP. It is demonstrated in [1] that

the mean-field equation of motion for the oneparticle density of the TASEP is equivalent to the exact Euler-Lagrange equations for the equilibrium density profiles of a binary mixture. Especially in this article, the aforementioned connection is to be applied for the PASEP. To our knowledge this relationship has not been published.

Classical density functional theory (DFT) [15, 16] is a powerful tool based on a variational principle which has become a standard framework to study static physical properties of equilibrium systems. The theory is applied to study various phenomena in inhomogeneous fluids, such as freezing and surface and interface behavior [17]. Classical DFT is recently used to treat aromatic ionic liquids at room temperature [18] and hydration of complex surfaces [19]. In principle, the aim of classical DFT is to construct an (approximate) free energy functional F of a classical system. Moreover, F can be written as a functional of the one-body density, $\rho(\mathbf{r})$, that is $F[\rho(\mathbf{r})]$, where \mathbf{r} is the position of an averaged number of particles. $F[\rho(\mathbf{r})]$ may eventually be determined via the Legendre transform of the grand potential, $\Omega[\rho(\mathbf{r})]$, which is also a functional of $\rho(\mathbf{r})$. Functional derivate of $F[\rho(\mathbf{r})]$ produces the one-body

direction correlation function, viz. $c^{(1)}$. Finally, minimizing $\Omega[\rho(\mathbf{r})]$ with respect to $\rho(\mathbf{r})$ yields the true one-body density at equilibrium. In this article, a method of classical DFT for hard core lattice models is utilized, i.e. the lattice fundamental measure theory (LFMT). LFMT is initially proposed in [20-22] as an extension of the FMT [23], which is developed based upon the molecular volume and other fundamental geometric measures of individual molecules [24].

THE FREE ENERGY FUNCTIONAL

The free energy functional only depends upon the interaction of the constituents of the system. The quantity is conveniently split into two parts, which is the ideal and the excess over ideal free energies, i.e.: $F_{\rm id}$ and $F_{\rm ex}$, respectively. For lattice systems the excess free energy functional in one-dimension (1D) may be defined as

$$F_{\text{ex}}\left[\rho_{i=1}(x), \dots \rho_{N}(x)\right] = F\left[\rho_{1}(x), \dots \rho_{N}(x)\right] - F_{\text{id}}\left[\rho_{1}(x), \dots \rho_{N}(x)\right], \tag{1}$$

where i = 1, ..., N denotes the species present in the system and $\rho_i(x)$ is the density of species i at lattice site x. The ideal free energy functional for lattice systems may be expressed as

$$F_{\rm id} = \frac{1}{\beta} \sum_{i=1}^{N} \sum_{x \in S} \rho_i(x) [\ln(\rho_i(x)) - 1], \tag{2}$$

with $1/\beta = k_B T$, k_B is the Boltzmann constant, T is the temperature, and S is the lattice system. The excess free energy functional for any 1D hard-core lattice fluids is

$$F_{ex}[\rho] = \frac{1}{\beta} \sum_{x \in S} \sum_{k \in I} a_k \Phi_0(n^{(k)}(x)), \tag{3}$$

where I is the set of indices suitably chosen to label every weighted densities $n^{(k)}(x)$, Φ_0 is the excess free energy of the 0D cavity [25], and a_k is the integer coefficient of Φ_0 . The weighted density $n^{(k)}(x)$ may be defined as

$$n^{(k)}(x) = \sum_{i=1}^{N} \sum_{t \in C_{k,i}^{(k)}} \rho_i(t), \tag{4}$$

where $C_{i,0D}^{(k)}$ is the k-th subset of $x \in S$, which is the appropriate 0D cavity of species i. Finally, the excess free energy functional for any 0D cavity in 1D is given by

$$\Phi_0(\eta) = (1 - \eta)\ln(1 - \eta) + \eta, \tag{5}$$

with $0 \le \eta \le 1$ is the averaged number of particles in the 0D cavity.

It may be observed that the ideal free energy functional is already provided by equation (2). However, the excess free energy functional is still unknown and depends upon the system under consideration. There is currently three methods to approximate functional (3), that is i) algebraic method, ii) diagrammatic method, and iii) Mobius inversion method. Here, the diagrammatic method is demonstrated. But before the method is used, as mentioned above, the models under consideration and their relationship have to be explained.

THE MODELS AND THEIR RELATIONSHIP

The dynamical model studied is the PASEP. This model consists of discrete lattice sites, S, and hard-core particles occupying the lattice sites. At each time steps, a site, $x \in S$, is chosen randomly with probability 1/N. If there is a particle on the chosen site, the particle may jump to the right (left)-nearest neighbor site with hopping rate k_+ (k_-) provided that there is no particle occupying the right (left)-nearest neighbor site. This is illustrated in Fig. 1.




FIGURE 1. The PASEP in 1D. The lattice sites are the vertical lines denoted by x = 1, 2, 3, ..., N. The big dark dots are the hard-core particles. The arrows show the movement of particles. The crosses show that the particles are not allowed to jump.

From the model above, the averaged quantities which we are interested in are the current density of particles moving to the right (left) and the density of particles occupying the sites, i.e.: $j_{+}(x)$ [$j_{-}(x)$] and $\rho(x)$, respectively. These quantities describe the macroscopic behavior of the PASEP. According to [1], the aforementioned quantities may be obtained via a connection to the appropriate equilibrium lattice fluids mixture. This connection is given as follows. A particle that jumps to the right (left)nearest neighbor site (see Fig. 1) corresponds to a dimer that has site exclusion and excludes its right (left)-nearest neighbor site (see Fig. 2), i.e. denoted by L₁ and L₂, respectively. A particle that does not jump (stays at the site) corresponds to a monomer (L₃) that only excludes its own site.




FIGURE 2. The equilibrium hard-core lattice fluids mixture with three species. L_1 and L_2 are dimers with positions at x = 2 and x = N - 1, respectively. L_3 is a monomer

From the correspondence above, the lattice fluids mixture with three species is obtained (Fig. 2). In this system, the averaged quantities are the densities of L_1 , L_2 , and L_3 , viz.: $\rho_{L_1}(x)$, $\rho_{L_2}(x)$, and $\rho_{L_3}(x)$, respectively. These three quantities are the Euler-Lagrange equations obtained later from the LFMT. Following the correspondence above, the relationship between the averaged quantities of the lattice fluids mixture and the PASEP is provided as follows. The densities of the dimers, i.e.: $\rho_{L1}(x)$ and $\rho_{L2}(x)$, may be realized as the density currents, i.e.: $j_+(x)$ and $j_-(x)$, respectively, whereas the monomer, $\rho_{L3}(x)$, may be realized as the density of the particles of the PASEP, $\rho(x)$. Hence, the task now is to determine $\rho_{L1}(x)$, $\rho_{L2}(x)$, and $\rho_{L3}(x)$ through the diagrammatic method of LFMT, such that $j_{+}(x)$, $j_{-}(x)$, and $\rho(x)$ may be finally gained.

F_{ex} CALCULATION VIA THE DIAGRAMMATIC METHOD

Before we determine the Euler-Lagrange equations, the $F_{\rm ex}$ has to be calculated in advance. Construction of the $F_{\rm ex}$ begins by determining the maximal cavities [25] of the species, i.e. \bigcirc ,

$$\bigcirc$$
 , and \bigcirc for L_1, L_2 , and L_3 , respectively.

The circles represent the lattice sites where the position of the particles is denoted by x. Hence, the maximal cavities of the mixture are

It may be observed that the maximal cavities above are the combination of the maximal cavities of the species. Hence, the first guess of the $F_{\rm ex}$ in accordance with equation (3) is attained as

$$\beta F_{\text{ex}} = \sum_{x \in S} \left[\Phi_0 () \right] + \Phi_0 ()$$

The functional (6) needs to be evaluated using any density profile, e.g.: \bigcirc as a test for the 0D

density profile. The evaluation produces spurious terms that have to be eliminated. This is done by introducing a second guess for the F_{ex} , i.e.:

$$\beta F_{\text{ex}} = \sum_{x \in S} \left[\Phi_0 \left(\bigotimes_x - \bigotimes_x \right) + \Phi_0 \left(\bigotimes_x - \bigotimes_x \right) \right] + \sum_{x \in S} \Phi_0 \left(\bigotimes_x - \bigotimes_x \right).$$

Using the same 0D density profile as a test for functional (7), yields the correct F_{ex} , that is

$$\beta F_{ex} = \sum_{x \in S} \left[\Phi_0 \left(\underbrace{\otimes}_{x} - \underbrace{\otimes}_{x} \right) + \Phi_0 \left(\underbrace{\otimes}_{x} - \underbrace{\otimes}_{x} \right) \right] + \sum_{x \in S} \left[\Phi_0 \left(\underbrace{\otimes}_{x} - \underbrace{\otimes}_{x} \right) + \Phi_0 \left(\underbrace{\otimes}_{x} \right) \right].$$
(8)

Moreover, since every cavity is associated to its density function, then we have from functional (8),

$$\beta F_{\text{ex}} = \sum_{x \in S} \left\{ \Phi_0 \left[\rho_{\text{L}_1}(x) + \rho_{\text{L}_1}(x+1) + \rho_{\text{L}_2}(x) + \rho_{\text{L}_3}(x) \right] + \Phi_0 \left[\rho_{\text{L}_1}(x-1) + \rho_{\text{L}_2}(x-1) + \rho_{\text{L}_2}(x) + \rho_{\text{L}_3}(x) \right] - \Phi_0 \left[\rho_{\text{L}_1}(x) + \rho_{\text{L}_2}(x+1) \right] - \Phi_0 \left[\rho_{\text{L}_1}(x) + \rho_{\text{L}_2}(x) + \rho_{\text{L}_3}(x) \right] \right\},$$
(2)

with each of the excess free energy of the 0D cavity, Φ, satisfies equation (5). Functional (9) is the exact excess free energy for the lattice fluids mixture with three species.

THE EULER-LAGRANGE EQUATIONS

The Euler-Lagrange equations can be determined using equations (1), (2), and (9). By calculating the direct correlation function for each species i, i.e.:

$$c_{\rho_i}^{(1)} = -\frac{\partial \beta F_{ex}}{\partial \rho_i(\mathbf{x}')},\tag{10}$$

then setting low densities of the dimers, that is, ρ_{L1} , $\rho_{L2} \rightarrow 0$, and finally inserting the results into the Euler-Lagrange equation, viz.:

$$\rho_i(x) = e^{c_{\rho_i}^{(1)}} e^{\beta V_i(x)}, \tag{11}$$

where V_i is any external potential acting upon species i, produces:

$$\rho_{L_1}(x) = e^{\beta [V_{L_1}(x) - V_{L_3}(x)]} \rho_{L_3}(x) [1 - \rho_{L_3}(x+1)], \quad (12)$$

$$\rho_{L_2}(x) = e^{\beta \left[V_{L_2}(x) - V_{L_3}(x) \right]} \rho_{L_3}(x) \left[1 - \rho_{L_3}(x-1) \right], \quad (13)$$

and

$$\rho_{L_3}(x) = e^{V_{L_3}(x)} (1 - \rho_{L_3}(x)). \tag{14}$$

Equations (12) - (14) are in fact the density profiles of the species in the equilibrium lattice fluids mixture system.

THE DENSITY AND CURRENT DENSITY OF THE PASEP

Finally, the macroscopic description of the PASEP, especially its one-particle and current densities may be achieved using equations (12) – (13). Invoking again the relationship given in the previous part, that is $j_+(x) \to \rho_{L1}(x)$, $j_-(x) \to \rho_{L2}(x)$, and $\rho(x) \to \rho_{L3}(x)$, and also introducing two more relationships, $e^{\beta \left[V_{L_1}(x)-V_{L_3}(x)\right]} \to k_+(x)$ and $e^{\beta \left[V_{L_2}(x)-V_{L_3}(x)\right]} \to k_-(x)$, produces

$$j_{+}(x) = k_{+}(x)\rho(x)[1 - \rho(x+1)], \tag{15}$$

and

$$j_{-}(x) = k_{-}(x)\rho(x)[1 - \rho(x - 1)]. \tag{16}$$

In addition, the density of the PASEP may be acquired by bringing into play the continuity equation,

$$\left. \frac{\partial \rho(x,t)}{\partial t} \right|_{\text{tot}} = -\left(\nabla j_{+} + \nabla j_{-} \right),\tag{17}$$

with $\nabla j_+(x) \equiv j_+(x-1) - j_+(x)$ and $\nabla j_-(x) \equiv j_-(x+1) - j_-(x)$. In steady state, the LHS of equation (17) goes to zero, which then produces

$$\rho(x) = \frac{\{k_{+}\rho(x-1)[1-\rho(x)] + k_{-}\rho(x+1)[1-\rho(x)]\}}{k_{+}[1-\rho(x+1)] + k_{-}[1-\rho(x-1)]},$$
(18)

which is just the density of the PASEP at steady state. It may be observed that equations (15), (16), and (18) are essentially mean-field equations. They are not the exact solution for the PASEP. However, the Euler-Langrange equations of the lattice fluids mixture are exact. Equation (17) is not a closed equation but may easily be solved self-consistently using numerical methods. It may also be noticed that setting one of the hopping rate to zero, e.g. $k \cdot (x) = 0$ in (16) yields back the known TASEP model.

CONCLUSION

The relationship between the PASEP and the lattice fluids mixtures has been presented. The relationship is based upon a correspondence between the particles and the jumping of particles of the PASEP with the species of the lattice fluids mixture. The (excess) free energy functional is calculated using the LFMT via a diagrammatic method, such that the Euler-Lagrange equations are obtained. Hence, the density and

current densities of the PASEP are achieved by invoking the aforementioned relationship.

ACKNOWLEDGMENTS

WSBD would like to thanks the Physics Education Department of Universitas Negeri Yogyakarta for the funding of this research.

REFERENCES

- W. S. B. Dwandaru and M. Schmidt, J. Phys. A: Math Theor. 40, 13209 (2007).
- L. Ciandrini, I. Neri, J. C. Walter, O. Dauloudet, and A. Parmeggiani, *Phys. Biol.* 11, 056006 (2014).
- D. Johann, D. Goswami, and K. Kruse, *Phys. Rev. E*. 89, 042713 (2014).
- E. Marshall, I. Stansfield, and M. C. Romano, J. R. Soc. Interface 11, 20140589 (2014).
- G. Poker, Y. Zarai, M. Margaliot, and T. Tuller, J. R. Soc. Interface 11, 20140713 (2014).
- Y. M. Yuan, R. Jiang, R. Wang, M. B. Hu, and Q. S. Wu, J. Phys. A: Math. Theor. 40 12351 (2007).
- R. Septiana, A. Indriawati, and W. S. B. Dwandaru, *Makara J. Sci.* 18(3), 79-85 (2014).
- 8. M. Zaim and W. S. B. Dwandaru, Jurnal Materi dan Pembelajaran Fisika, 4(2), 38-45 (2014).
- H. Ito and K. Nishinari, Phys. Rev. E 89, 042813 (2014).
- D. Chowdhury, K. Nishinari, and A. Schadschneider, Phase Transition 77, 601-624 (2004).
- D. Chowdhury, A. Schadschneider, K. Nishinari, *Physics of Life Reviews* 2, 318-352 (2005).
- 12. S. Srinivasa and M. Haenggi, The TASEP: A Statistical Mechanics Tool to Study the Performance of Wireless Line Network, in the Proceeding of the 19th IEEE ICCCN 2010, Zurich, Switzerland, 2010.
- C. Landim, S. Olla, and S. R. S. Varadhan, Commun. Math Phys 224, 307-321 (2001).
- W. S. B. Dwandaru, "Various Correspondences Between Simple Driven and Equilibrium Statistical Hard Core Models", Ph.D. Thesis, Bristol University, UK, 2010.
- P. J. Hasnip, K. Refson, M. I. J. Probert, J. R. Yates, S. J. Clark, and C. J. Pickard, *Phil. Trans. R. Soc. A* 372 20130270 (2014).
- 16. R. Evans, Adv. Phys. 28, 143 (1979).
- M. Schmidt, J. Phys.: Condens. Matter 15, S101 (2003).
- M. Turesson, R. Szparaga, K. Ma, C. E. Woodward, and J. Forsman, *Soft Matter* 10, 3229-3237 (2014).
- G. Jeanmairet, V. Marry, M. Levesque, B. Rotenberg, and D. Borgis, *Molecular Physics* 112, 1320-1329 (2014).
- L. Lafuenta and J. A. Cuesta, *Phys. Rev. Lett.* 93, 130603 (2004).
- L. Lafuente and J. A. Cuesta, J. Phys. A: Math. Gen 38, 7461 (2005).

THE 4th INTERNATIONAL CONFERENCE ON THEORETICAL AND APPLIED PHYSICS (ICTAP-2014) 16-17 October 2014, Denpasar-Bali, Indonesia

- L. Lafuente, J. A. Cuesta, and M. Schmidt, *Phys. Rev.* E 72, 0331405 (2005).
- 23. Y. Rosenfeld, *Phys. Rev. Lett.* **63**, 980 (1989).
- P. Tarazona and Y. Rosenfeld, *Phys. Rev. E* 8, L577 (1996).
- W. S. B. Dwandaru, "Theory of Dynamics in Driven Lattice Fluids", Master by Research Dissertation, Bristol University, UK, 2006.




ISBN: 978-602-294-116-3 | Published: 12 May 2016



PROCEEDING THE 4th INTERNATIONAL CONFERENCE ON THEORETICAL AND APPLIED PHYSICS (ICTAP) 2014

Bali, Indonesia 16 - 17 October 2014



Editors:

Ni Nyoman Rupiasih, Wayan Gede Suharta and Hery Suyanto Udayana University, Bali, Indonesia

All Papers have been peer reviewed.

Copyright © 2016, by Udayana University Press. All rights reserved.

Printed by: Udayana University Press, Jl. P.B. Sudirman, Denpasar, Bali, Indonesia 80114

ISBN: 978-602-294-116-3 Published: 12 May 2016

PROCEEDING THE 4th INTERNATIONAL CONFERENCE ON THEORETICAL AND APPLIED PHYSICS (ICTAP) 2014

Bali, Indonesia 16-17 October 2014

Editors:

Ni Nyoman Rupiasih, Wayan Gede Suharta and Hery Suyanto Udayana University, Bali, Indonesia

Cover Design: I Ketut Putra Udayana University, Bali, Indonesia

Sponsoring Organizations Indonesian Physical Society – IPS Indonesian Physical Society – Bali Branch

All papers have been peer reviewed

E-411-PE-504-87P NZ

Copyright © 2016 by Udayana University Press. All rights reserved. Printed by: Udayana University Press, Jl. P.B. Sudirman, Denpasar, Bali, Indonesia 80114

PREFACE

The 4th International Conference on Theoretical and Applied Physics (ICTAP) 2014 was held in Bali, Indonesia on October 16-17, 2014. It is an international conference covering a wide subject in the field of theoretical and applied physics. This Conference was organized by the Department of Physics, Udayana University (UNUD) and Indonesian Physical Society (HFI). Many people have unreservedly and enthusiastically participated in the planning and preparation of this conference, including the Organizing and Scientific Committees, the speakers, the conference officer, students, etc. Thank you all for your positive attitude and fantastic support. Thanks to all invited speakers: Prof. Dr. Yusaku Fujii (Gunma University, Japan), Prof. Jakrapong Kaewkhao, Ph.D (Nakhon Pathom Rajabhat University, Thailand), Prof. Dr. Hong Joo Kim (Kyungpook National University, Republic of Korea) and Prof. Dr. Ing Mitra Djamal (ITB) that you have accepted our invitation. The financial support from Udayana University is gratefully acknowledged. ICTAP 2014 is aimed at providing the much needed forum of scientific communication and interaction among distinguished scientists working in the field of physics and related fields.

This event was also designed to offer the opportunity of making direct contact for the young Indonesian scientists and students with well-known scientists abroad and thereby fostering the existing research collaborations and extending international research networking for the future.

More than 106 authors from 6 countries have submitted their work in the conference. ICTAP 2014 finally accepted 57 original research papers after a peer review process. During the conference, 14 parallel sessions were held in order to advance and contribute to specific research area in physics.

Finally, special thanks to you, the delegates, for supplying the input needed for successful scientific conferences. We wholeheartedly welcome you and hope you find ICTAP 2014 as successful and rewarding as we envision it to be.

Denpasar, 12 May 2016



Ni Nyoman Rupiasih, Ph.D. Dr. Wayan Gede Suharta Dr. Hery Suyanto

Table of Contents

| | PLENARY AND INVITED PAPERS | page |
|---------|--|---------|
| KS-1 | Review of the Levitation Mass Method (LMM) – A Precision Method for Measuring Mechanical Quantities Using An Optical Interferometer, <i>Yusaku Fujii</i> , <i>Hadi Nasbey</i> , <i>Agus Setyobudi and Akihiro Takita</i> | |
| KS-2 | White Emission Materials from Glass Doped with Rare Earth Ions, <i>P. Yasaka and J. Kaewkhao</i> | 007-020 |
| | Parallel Session TC: Theoretical Physics and Computation | |
| TC 1-01 | Application of quantum heat engineon the three-dimensional potential, <i>Andika Kusuma Wijaya</i> | 021-026 |
| TC 1-05 | Direct Method of Calculus of Variation on Electromagnetism in Differential Form, I Gusti Ngurah Yudi Handayana and Muhammad Farchani Rosyid | 027-034 |
| TC 1-07 | Chaos Identification of the Double Pendulum Motion in the Magnetic Field by Joko Saefan and Suparmi | 035-040 |
| TC 1-10 | Computer Simulation Development base on Open Source to improve Students Conceptual Competence, <i>Madlazim and Dyah Permatasari</i> | 041-050 |
| TC 1-15 | Quantum Physics Learning by Using Matlab 7.6 on Highlights Potential Transmission Coefficient Parabolic One Dimension, <i>Reni Herniati and Andika Kusuma Wijaya</i> | 051-055 |
| TC 1-21 | A Connection for a Partially Asymmetric Driven Lattice Gas of A Mean-Field Type to an Exact Lattice Density Functional Via a Diagrammatic Method, Wipsar Sunu Brams Dwandaru and Matthias Schmidt | 056-061 |
| TC 1-23 | Theoretical EQUATION WITH EVIDENCE Relaxation oscillator using an oscilloscope ON THE CIRCUITS OP-AMP, <i>Yohanes Soenarto</i> , <i>Imas Ratna Ermawaty</i> , <i>Tri Isti Hartini and Felicianda</i> | 062-069 |
| | Parallel Session AM: Advanced Material and Nanotechnology | |
| AM 1-02 | LPG Gas Sensing Properties of Composite CuO(TiO2) Sensor, Elvaswer, Muhammad Faisal, Dwi Puryanti, Essy Puspa Zelvia and Ratna Sari Dewi | 070-073 |
| AM 1-04 | Electrodeposition and Characterization of Ni-TiAlN Composite Film, Esmar Budi, Agus Setyo Budi, Iwan Sugihartono, Setia Budi and Hadi Nasbey | 074-079 |
| AM 1-06 | Effect of Cold Isostatic Pressing on Properties of Millimeter Wave Sintered High Purity Alumina, <i>Ida Usman, I Nyoman Sudiana, La Ode Ngkoimani and Usman Rianse</i> | 080-083 |
| AM 1-07 | The Preparation of Natural Rubber –g-Glysidil Metacrilate Nanocomposite, Kurnia Sembiring and Riani Sari Sembiring | 084-092 |
| AM 1-29 | Characterization Of ZnO Thin Films Doped with Natrium by Sol-Gel Method, P.L. Gareso, N. Syuhada, N. Rauf, E. Juarlin, Sugianto and A. Maddu | 093-097 |

| AM 1-30 | The Influence of the Milling Time Process on Microstructure and Magnetic Properties of Isotropic Bonded NdFeB Magnets, <i>Priyo Sardjono</i> , <i>Muljadi</i> , <i>Nenen Rusnaeni</i> , <i>Suprapedi and Ayu Yuswitasari</i> | 098-10 |
|---------|---|--------|
| | Parallel Session BM: Biophysics and Medical Physics | |
| BM 1-01 | Concentrations of Some Natural Occurring Radionuclides and Particles during the 2010 Eruption of Mount Bromo in East Java, Indonesia, <i>Johan A.E. Noor</i> | 103-10 |
| BM 1-02 | Automatic thresholding with Otsu Method To Identify Plasmodium falciparum Phase in Malaria-infected Red Blood Cells, <i>Kusworo Adi</i> , <i>Sri Pujiyanto</i> , <i>Rahmat Gernowo</i> , <i>Adi Pamungkas and Ari Bawono Putranto</i> | 107-1 |
| BM 1-04 | Head Impact Analysis of Children in Soccer Heading, Nugroho Agung S., Tommy Apriantono and Suprijanto | 113-1 |
| BM 1-06 | Development of Low-Cost Modular Wireless EEG System, Andri Rahmadhani, Sra H. Pratama, Suprijadi, Freddy Haryanto and Suparno Satira | 118-1 |
| BM 1-08 | Analyzis of Computed Tomography Dose Index (CTDI) value towards X-ray Tube Current and Voltage Variations of Computed Tomography Scanner (CT Scan) by using PPMA Phantom, Syamsir Dewang, Bualkar Abdullah, Bannu, Nur Hasanah, Suryaningsih and Satrial Male | 123-1 |
| BM 1-11 | Burnable Poison Neutronic Characteristics of Hexagonal Tight Lattice Cell for Small Long-Life BWR with Thorium Based Fuel, Nuri Trianti, Zaki Su'ud, Idam Arif, Sidik Permana and Eka Sapta Riyana | 128-1 |
| | Parallel Session G: Geophysics | |
| G 1-01 | Estimation of Ore Body Distribution at X-Field, Nangroe Aceh Darussalam Province, Using Resistivity Method, Dipole-Dipole Configuration, Adi Susilo | 135-1 |
| G 1-02 | Estimating Reservoar Temperature Using Geothermometer Equation On Hot Springs In Panti District, Pasaman Regency, West Sumatera, Indonesia, <i>Ardian Putra and Rahmat Arrahman</i> | 140-1 |
| G 1-03 | Relation Model between Electrical and Mechanical Properties on the Exploration Sub Surface under Ground, Lantul, D.A. Suriamihardja, A.M. Imran and Tri Harianto | 144-1 |
| G 1-04 | The Geothermal Model of Dieng Plateau Complex from Resistivity Image With Magnetotelluric Method, <i>Eddy Z Gaffar</i> | 148-1 |
| G 1-07 | Application of Geoelectrical Resistivity Method to Investigate Subsurface Geology Structure of Lava Spill Area of Gamalama Volcano, <i>Fatma Hamid, Saprudin, M. Toifur and Yudhiakto Pramudya</i> | 154-1 |
| G 1-08 | Analysis Of Subsurface Materials Based On The Price Of Medium Permeability In The Karst Region Pangkep, <i>Muhammad Arsyad</i> , <i>Nasrul Ihsan and Vistarani Arini Tiwow</i> | 160-1 |
| G 1-17 | The Analysis of P-Wave Period-Duration as One of Parameters for Tsunami Early Warning, Zulkarnain Adnan, Sugeng Pribadi and Nanang T. Puspito | 165-1 |
| G 1-18 | 10 | 171-1 |

| | Parallel Session A: Astrophysics | |
|---------|---|---------|
| A 1-01 | Study of Ideal Magnetohydrodynamics In Curved Space, Luh Putu Budi Yasmini, and I Gede Aris Gunadi | 175-178 |
| | Parallel Session IP: Instrumental Physics | |
| IP 1-01 | Development of Web-based Power Monitoring System for a Pulse Combustion Spray Pyrolysis using Java Programming, Agus Fattrya Nanda, Darmawan Hidayat, Camellia Panatarani, Dwindra W. Maulana and I Made Joni | 179-183 |
| IP 1-03 | Digital Receiver Synchronization Methods for OFDM Modem, Assa'idah and Hadi | 184-189 |
| | Parallel Session LO: Laser and Opto-electronics | |
| LO 1-02 | Dependence of the defect modes on the strength of localized defects in photonic lattice, Arif Hidayat, Diana Kurniati, Hari Wisodo and Eny Latifah | 190-197 |
| LO 1-05 | Potential Candidate for Gigahertz range Electromagnetic wave absorbers of Carbon Sheets Based on Polymer, Maria Margaretha Suliyanti, Nanik Indayaningsih and Affi Nurhidayah | 198-203 |
| LO 1-08 | Study on the Theory of Electron Multiple Scattering on Plasma, <i>Taat Guswantoro</i> , <i>Muhammad Nur and Vincencius Gunawan</i> | 204-209 |
| | Parallel Session EE: Energy and Environment Physics | |
| EE 1-03 | Development Of Wind Tunnel For Aerodynamic Test To Support The Wind Energy Research, <i>Hadi Nasbey</i> , <i>Cecep Rustana and Christine Stefphanie</i> | 210-215 |
| | Parallel Session I: Interdisciplinary Physics | |
| I 1-01 | A Misunderstanding of Force Concept; the action-reaction pairs same as a cause-effect sequence, A. Halim, Melvina and Susilawati | 216-219 |
| I 1-04 | A Study on The Physical Characteristics of Lemukutan Island Territorial Water Wave in Bengkayang Regency, Eka Murdani and Sumarli | 220-224 |
| I 1-08 | The Influence of Cooperative Instructional Strategies with STAD Technique on Students Achievement in Physics of Grade XI SMA Lab School, Palu, Marungkil Pasaribu | 225-229 |
| I 1-11 | The Relationship between Formal Thinking Abilities and Problem-Solving Skills in Kinematics Topics, Sondang R. Manurung | 230-234 |
| I 1-12 | Theoretical Modeling for the Effect Tenacity on Take-Up Roller (Ro) and Tenacity on Winding Device (Rw) Related With the Yarn Breakage on Rotor Open End | 235-240 |
| | Spinning, Valentinus Galih Vidia Putra, M.F. Rosyid and R. Arief Dewanto | |

${\tt C62_Proceeding_ICTAP_A_Connection_for_A_Partially.pdf}$

| ORIGINA | LITY REPORT | 0 | | - 71 |
|-------------|---|--|--|----------------------|
| 2 SIMILA | % RITY INDEX | 20% INTERNET SOURCES | 18% PUBLICATIONS | 3% STUDENT PAPERS |
| PRIMAR | Y SOURCES | | | |
| 1 | aip.scita Internet Source | tion.org | | 2% |
| 2 | www.fis | ikanet.lipi.go.ic | | 2% |
| 3 | orbit.dtu Internet Source | | | 1 % |
| 4 | www.thp | ohy.uni-duesse | ldorf.de | 1 % |
| 5 | lipi.go.io | | | 1 % |
| 6 | Submitte Student Paper | ed to Universit | as Brawijaya | 1 % |
| 7 | Darmaw Couloml totally a with a si | van. "Towards to b blockade in c symmetric sim | vandaru, Denny the dynamics of quantum dot via ple exclusion p anced Studies i 16 | f |

| _ | 8 | "Strategic Innovative Marketing", Springer Science and Business Media LLC, 2017 Publication | 1 | % |
|---|----|--|---|---|
| | 9 | journal.uny.ac.id Internet Source | 1 | % |
| _ | 10 | repository.unhas.ac.id Internet Source | 1 | % |
| | 11 | "Preface", Journal of Physics: Conference Series, 2021 Publication | 1 | % |
| | 12 | Corinne Aubert, Olivier Buisine, M. Petit, F. Slowinski, Max Malacria. "Cobalt-mediated cyclotrimerization and cycloisomerization reactions. Synthetic applications", Pure and Applied Chemistry, 1999 Publication | 1 | % |
| _ | 13 | sintadev.ristekdikti.go.id Internet Source | 1 | % |
| _ | 14 | www.unud.ac.id Internet Source | 1 | % |
| _ | 15 | mafiadoc.com Internet Source | 1 | % |
| | 16 | Geoffroy, M.H "Local convergence of some iterative methods for generalized equations", Journal of Mathematical Analysis and Applications, 20040215 | 1 | % |

| 17 | Róbert Juhász. "Weakly coupled, antiparallel, totally asymmetric simple exclusion processes", Physical Review E, 2007 Publication | 1 % |
|----|---|-----|
| 18 | "Traffic and Granular Flow '07", Springer Science and Business Media LLC, 2009 | <1% |
| 19 | psp.ub.ac.id Internet Source | <1% |
| 20 | www.citeulike.org Internet Source | <1% |
| 21 | www.scientific.net Internet Source | <1% |
| 22 | Izzati Muhimmah, Nani Harniawati, Novyan Lusiyana. "Characteristics determination of infected erithrocytes by plasmodium falciparum as a diagnostic of malaria, based on microscopic images", 2017 International Conference on Advances in Computing, Communications and Informatics (ICACCI), 2017 Publication | <1% |
| 23 | Wipsar Sunu Brams Dwandaru, Matthias Schmidt "Variational principle of classical | <1% |

Schmidt. "Variational principle of classical


density functional theory via Levy's

constrained search method", Physical Review E, 2011

Publication

| 24 | pt.scribd.com Internet Source | <1% |
|----|---|-----|
| 25 | www.coursehero.com Internet Source | <1% |
| 26 | Astri Novianty, Carmadi Machbub, Sri Widiyantoro, Irwan Meilano, Hendy Irawan. "Tsunami Potential Identification based on Seismic Features using KNN Algorithm", 2019 IEEE 7th Conference on Systems, Process and Control (ICSPC), 2019 Publication | <1% |
| 27 | www.magistergeophys.sci.unhas.ac.id | <1% |
| 28 | digilib.unimed.ac.id Internet Source | <1% |
| 29 | fmipa.itb.ac.id Internet Source | <1% |
| 30 | E. Marshall, I. Stansfield, M. C. Romano. "Ribosome recycling induces optimal translation rate at low ribosomal availability", Journal of The Royal Society Interface, 2014 Publication | <1% |
| | | |

| <1 % <1 % |
|-------------------------|
| /1 _{0/} |
| \ % |
| <1% |
| <1% |
| <1% |
| <1% |
| <1% |
| |



| 45 | arxiv.org Internet Source | <1% |
|----|---|------|
| 46 | e-archivo.uc3m.es Internet Source | <1% |
| 47 | www.mipa.sci.unhas.ac.id Internet Source | <1 % |
| 48 | W S Brams Dwandaru, Matthias Schmidt. "A relationship of mean-field theory for a driven lattice gas to an exact equilibrium density functional", Journal of Physics A: Mathematical and Theoretical, 2007 Publication | <1% |
| 49 | A. Mani. "Chapter 9 Esoteric Rough Set Theory: Algebraic Semantics of a Generalized VPRS and VPFRS", Springer Science and Business Media LLC, 2008 Publication | <1 % |
| 50 | Luis Lafuente. "Cluster density functional theory for lattice models based on the theory of Möbius functions", Journal of Physics A Mathematical and General, 08/26/2005 | <1% |
| | 1 diffication | |

Case", IEEE/ACM Transactions on

Computational Biology and Bioinformatics, 2014

Publication



Yuri Martínez-Ratón, José A. Capitán, José A. Cuesta. "Fundamental-measure density functional for mixtures of parallel hard cylinders", Physical Review E, 2008
Publication

<1%



www.sciepub.com

Internet Source

<1%

Exclude quotes Off
Exclude bibliography On

Exclude matches

Off

${\tt C62_Proceeding_ICTAP_A_Connection_for_A_Partially.pdf}$

GRADEMARK REPORT

FINAL GRADE

GENERAL COMMENTS

/100

Instructor

| PAGE 1 | |
|---------|--|
| PAGE 2 | |
| PAGE 3 | |
| PAGE 4 | |
| PAGE 5 | |
| PAGE 6 | |
| PAGE 7 | |
| PAGE 8 | |
| PAGE 9 | |
| PAGE 10 | |
| PAGE 11 | |
| PAGE 12 | |