

MEETING: 17-18

Learning Objectives: Students are able to

1. explain analytic functions
2. explain harmonic functions
3. determine a harmonic conjugate

1. Analytic Functions

Definition

- (a). A function f of the complex variable z is *analytic* at point z_0 if its derivative exists not only at z_0 but at each point z in some neighborhood of z_0 .
- (b). An *entire* function is a function that is analytic at each point in the entire plane.
- (c). If a function fails to be analytic at a point z_0 but is analytic at some point in every neighborhood of z_0 is called a *singular point* or *singularity of the function*.

Example: the function

$$f(z) = \frac{1}{z} \text{ then } f'(z) = -\frac{1}{z^2}$$

We can see that f is analytic except in the point $z = 0$ in the finite plane. Moreover, $z = 0$ is called a singular point.

2. Harmonic Functions

A real-valued function h of two real variables x and y is said to be *harmonic* in a given domain of the xy plane if, throughout that domain, it has continuous partial derivatives of the first and second order and satisfies the partial differential equation

$$h_{xx}(x, y) + h_{yy}(x, y) = 0 \tag{21}$$

known as Laplace's equation.

Example

The function $H(x, y) = e^{-y} \sin x$ is harmonic since $H_x = e^{-y} \cos x$, $H_{xx} = -e^{-y} \sin x$, $H_y = -e^{-y} \sin x$, $H_{yy} = e^{-y} \sin x$ and $H_{xx}(x, y) + H_{yy}(x, y) = 0$

3. Harmonic Conjugate

If two given functions u and v are harmonic in a domain D and their first partial derivatives satisfy the Cauchy-Riemann equations throughout D , we say that v is a *harmonic conjugate* of u . We now illustrate one method of obtaining a harmonic conjugate of a given harmonic function.

The function

$$u(x, y) = y^3 - 3x^2y.$$

To find a harmonic conjugate $v(x, y)$, we note that $u_x(x, y) = -6xy$.

So, in view of the condition $u_x = v_y$ we may write $v_y = -6xy$.

Holding x fixed and integrating both sides of this equation with respect to y , we find that

$$v(x, y) = -3xy^2 + g(x)$$

where $g(x)$ is arbitrary function of x . Since the condition $u_y = -v_x$ must hold, it follows that

$$3y^2 - 3x^2 = 3y^2 - g'(x)$$

So, $g'(x) = 3x^2$, and this means that $g(x) = x^3 + c$, where c is an arbitrary real number. Hence the function

$$v(x, y) = x^3 - 3xy^2 + c$$

is a harmonic conjugate of our function. The corresponding analytic function is

$$f(z) = (y^3 - 3x^2y) + i(x^3 - 3xy^2 + c)$$

It is easily verified that

$$f(z) = i(z^3 + c)$$

Exercises:

1. In each case, determine the singular points of the functions and state why the function is analytic everywhere except at those points

a. $f(z) = \frac{z+1}{z(z^2+2)}$ b. $f(z) = \frac{z}{z^2-2iz+3}$

2. Show that the following functions are entire

a. $f(z) = \cosh x \sin y - i \sinh x \cos y$

b. $g(z) = \cosh x \cos y + i \sinh x \sin y$

3. Let a , b and c be real constants. Determine a relation among the coefficients that will guarantee that the function $f(x, y) = ax^2 + bxy + cy^2$ is harmonic.

4. Show that $u(x, y)$ is harmonic in some domain and find a harmonic conjugate $v(x, y)$ when

a. $u(x, y) = 2x(1 - y)$

b. $u(x, y) = 2x - x^3 + 3xy^2$

5. Let $u_1(x, y) = x^2 - y^2$ and $u_2(x, y) = x^3 - 3xy^2$. Show that u_1 and u_2 are harmonic functions and that their product $u_1(x, y) \cdot u_2(x, y)$ is not a harmonic function.