

## 1. Derivatives

Let  $f$  be a function whose domain of definition contains a neighborhood of a point  $z_0$ . The *derivative* of  $f$  at  $z_0$ , written  $f'(z_0)$ , is defined by the equation

$$f'(z_0) = \lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0}, \quad (13)$$

provided this limit exist.

By expressing the variable  $z$  in definition (13) in terms of the new complex variable  $\Delta z = z - z_0$ , we can write that definition as

$$f'(z_0) = \lim_{\Delta z \rightarrow 0} \frac{f(z_0 + \Delta z) - f(z_0)}{\Delta z}. \quad (14)$$

Note that, because  $f$  is defined throughout a neighborhood of  $z_0$ , the number  $f(z_0 + \Delta z)$  is always defined for  $|\Delta z|$  sufficiently small. Let  $\Delta w = f(z_0 + \Delta z) - f(z_0)$ , then if we write  $\frac{dw}{dz}$  for  $f'(z)$ , equation (14) becomes

$$\frac{dw}{dz} = \lim_{\Delta z \rightarrow 0} \frac{\Delta w}{\Delta z}. \quad (15)$$

*Example*

Suppose that  $f(z) = z^3$ . At any point  $z$ ,

$$\lim_{\Delta z \rightarrow 0} \frac{\Delta w}{\Delta z} = \lim_{\Delta z \rightarrow 0} \frac{(z + \Delta z)^3 - z^3}{\Delta z} = \lim_{\Delta z \rightarrow 0} 3z^2 + 3z(\Delta z)^2 + (\Delta z)^2 = 3z^2. \text{ Hence } \frac{dw}{dz} = 3z^2 \text{ or } f'(z) = 3z^2.$$

## 2. Differentiation Formulas

Let  $c$  be a complex constant, and let  $f$  be a function whose derivative exist at a point  $z$ . It is easy to show that

$$\frac{d}{dz} c = 0, \quad \frac{d}{dz} z = 1, \quad \frac{d}{dz} cf(z) = cf'(z). \quad (16)$$

Also, if  $n$  is a positive integer

$$\frac{d}{dz} z^n = nz^{n-1} \quad (17)$$

### **Theorem**

If the derivation of two function  $f$  and  $g$  exist at a point  $z$ , then

$$(1) \frac{d}{dz} [f(z) + g(z)] = f'(z) + g'(z)$$

$$(2) \frac{d}{dz} [f(z)g(z)] = f'(z)g'(z)$$

$$(3) \frac{d}{dz} \left[ \frac{f(z)}{g(z)} \right] = \frac{f'(z)g(z) - g'(z)f(z)}{(g(z))^2}, \text{ when } g(z) \neq 0.$$

### **Example**

To find the derivative of  $(2z^2 + i)^5$ . According to the theorem, we have

$$\frac{d(2z^2 + i)^5}{dz} = 5(2z^2 + i)^4 (4z) = 20z(2z^2 + i)^4.$$

### **Exercises**

1. Apply definition (15) of derivative to find  $f'(z)$  when

a.  $f(z) = \frac{1}{z}, z \neq 0$

b.  $f(z) = 3z^2 - 2z$

c.  $f(z) = \left(z + \frac{1}{2}\right)^4$

2. Apply definition (15) of derivative to find  $f'(z)$  when  $f(z) = z^3 - 4z$  at point

a.  $z = z_0$

b.  $z = i$

3. Use result in Sec. 2 to find  $f'(z)$  when

a.  $f(z) = 3z^2 - 4z + 1$

b.  $f(z) = (1 - 3z^3)^2$

c.  $f(z) = \frac{z+i}{3z-2}$

4. If the derivation of two function  $f$  and  $g$  exist at a point  $z$ , proof that

$$\frac{d}{dz}[f(z) + g(z)] = f'(z) + g'(z).$$

5. Show that  $f'(z)$  does not exist at any point  $z$  when  $f(z) = \bar{z}$ .